

568-90
389548
N91P14168

NARROW BAND IMAGES OF SUSPECTED COOLING FLOW GALAXIES

Susana Deustua and Greg Bothun
Department of Astronomy
University of Michigan
Ann Arbor, MI 48109

ABSTRACT: We present narrow-band (H α and [NII]) CCD images of elliptical galaxies suspected of having cooling flows.

I. INTRODUCTION: Cooling flows analogous to those in clusters of galaxies are also present on smaller scales in isolated (i.e. not in the center of a cluster) early-type galaxies as evidenced by observed soft x-ray halo emission around these galaxies (Forman, Jones and Tucker (1985); Canizares, Fabbiano and Trinchieri (1987); Fabbiano et al (1987)). The typical luminosity in these halos is $L_x = 10^{39} - 10^{41}$ ergs s $^{-1}$. This gas most probably originates as mass loss from stellar evolution which has been driven out of the galaxy by supernovae induced winds and subsequently flows inward in a cooling flow (Mathews and Loewenstein 1986). These smaller flows should exhibit temperature and density gradients, optically emitting filaments and star formation (at the rate of up to a few M_\odot yr $^{-1}$). Therefore one expects to find cool gas ($T = 10^4$ K) in the inner few kpc of elliptical galaxies.

II. OBSERVATIONS: Narrow-band (20 Å bandpass) interference filter images in H α and [NII] of a subsample of 12 of approximately 40 cooling flow galaxies were taken at the MDM observatory with the 1.3m McGraw-Hill Telescope. Previous narrow-band imaging has been at a resolution insufficient to separate H α from [NII]. Typical exposure times are 900 seconds.

III. RESULTS and DISCUSSION: We find evidence of optical emission from cool gas which is most likely associated with a cooling flow. At least 4 galaxies (NGC3998, NGC4203, NGC4550 and NGC4697) show emission of [NII] and/or H α within the central 20", a fraction consistent with results from Phillips et al (1987) and Caldwell (1984). In these four, [NII] λ 6584Å emission is stronger than H α (which is often in absorption e.g. as seen in long slit spectra of NGC4697). From long slit spectra of these galaxies, Deustua and Teske (1989) infer electron densities, N_e , of order 1.2×10^3 cm $^{-3}$ assuming $T = 10^4$ K for the optically emitting gas. NGC 2685, NGC 3489 appear to have emission in [NII]; NGC 4636 may also, but, is difficult to see in our data (Demoulin-Ulrich, Butcher and Boksenberg (1984) did not see emission from this galaxy). NGC 4472, NGC 4473, NGC 4365, NGC 4638 and NGC 4649 show no emission.

93121-103

Table 1

Parameters of the Sample Galaxies

OBJECT	type	$\log L_x$ erg/s	a_x kpc	$N_e(0)$ cm^{-3}	$t_{cl}(0)$ 10^7 yr	M_{gas} $10^{10} M_\odot$
NGC 2685	S0	39.91	0.25	0.14	4.7	0.017
NGC 3489	S0/Sa	<39.80	0.24	0.13	5.2	0.015
NGC 3998	S0	41.74	0.44	0.50	1.4	0.338
NGC 4203	S0	41.34	0.40	0.36	1.9	0.186
NGC 4365	E3	40.29	0.65	0.05	13.2	0.117
NGC 4472	E1/S0	41.71	1.80	0.06	11.8	2.741
NGC 4473	E5	39.95	0.46	0.06	11.5	0.047
NGC 4550	E7/S0	<39.85	0.18	0.21	3.2	0.010
NGC 4636	E0/S0	41.64	0.70	0.22	3.1	0.618
NGC 4638	S0	39.59	0.22	0.11	5.9	0.011
NGC 4649	E0	41.40	1.16	0.08	8.7	0.988
NGC 4697	E6	40.27	0.95	0.03	23.4	0.199

Galaxy type is from the Revised Shapley-Ames Catalog. $\log L_x$ is given in Canizares, Fabbiano and Trinchieri (1987); the values of the core radius, a_x , central density, $N_e(0)$, central cooling time, $t_{cl}(0)$, and gas mass, M_{gas} , for the hot gas are calculated using their equations.

The best candidates for active cooling flows are NGC 3998 and NGC 4203. Both have H α and [NII] emission as well as large central densities and short central cooling times as calculated from their x-ray properties (Table 1), furthermore, van Driel *et al* (1988) observed an HI disk in NGC 4203. Whereas the galaxies with weaker [NII] emission and no H α have lower central densities and longer cooling times.

It has been suggested in the literature that distorted isophotes i.e. pointy or boxy, indicate the presence of accretion processes. Jedrzejewski (1987) found pointy isophotes for NGC 4473 and NGC 4697, and Mollenhoff and Bender (1986) found boxy isophotes in NGC 4365 and NGC 4472. However, while NGC 4697 has optical emission, its central $N_{e\text{Xray}}$ is small and t_{coolXray} is large. We find that broad-band U, B, and I images of 7 southern cooling-flow galaxies show that 3 have boxy isophotes at intermeditate radii and pointy isophotes at the inner and outermost radii (i.e. mixed), two are essentially boxy, one has elliptical isophotes and one, PKS0745-191, is too small to fit. Three of the control galaxies

1 00131-101

have isophotes which are essentially elliptical. However, IC 3370, while not a cooling flow candidate, also has very obvious boxy isophotes (Jarvis 1987).

Distorted isophotes probably confirm the presence of a cooling flow via its dynamical effect but do not alone provide sufficient evidence. Whether the presence of H α and [NII] emission alone are related to cooling flows is unclear (see also Donahue, Stocke and Voit this conference), though Hu, Cowie and Wang (1985) observed such emission filaments in several cluster cooling flows.

Acknowledgments : S. D. thanks the Astronomy Dept. at CWRU for use of their computing facilities, G. Aldering and R. E. Luck for useful suggestions. We also thank I. Thompson for obtaining some broad band images. Partial support for this research was provided by a Rackham Dissertation Grant from the University of Michigan.

REFERENCES:

- Caldwell, N., 1984 P. A. S. P., 96, 287
Canizares, C., Fabbiano, G. and Trinchieri, G., 1987 Ap. J., 312, 503
Demoulin-Ulrich, M-H., Butcher, H. R. and Boksenberg, A., 1984 Ap. J., 285, 527
Deustua, S. and Teske, R., 1989 in preparation
Fabbiano, G., Klein, U., Trinchieri, G. and Wielebinski, R., 1987 Ap. J., 312, 111
Forman, W., Jones, C. and Tucker, W., 1985 Ap. J., 293, 102
Hu, E., Cowie, L. and Wang, Z., 1985 Ap. J. Suppl., 59, 447
Jarvis, B., 1987 A. J., 94, 30
Jedrzejewski, R. I., 1987 M. N. R. A. S., 226, 747
Mathews, W. G. and Loewenstein, M., 1986 Ap. J. (Lett.), 306, L7
Mollenhoff, C., and Bender, R., 1986 in IAU Symposium 127: Structure and Dynamics of Elliptical Galaxies ed. T. de Zeew (Dordrecht: Reidel) p. 409
Phillips, M., Jenkins, C. R., Dopita, M. A., Sadler, E. M. and Binette, L., 1986 A. J., 91, 1062
van Driel, W., van Woerden, H., Gallagher III, J. S. and Schwarz, U. J., 1988 Astronomy and Astrophysics, 191, 201
Vedder, P. 1988 MIT thesis